

One important use of the preceding theorem is as a tool for verifying that a map is a homeomorphism:

**Theorem 26.6.** *Let  $f : X \rightarrow Y$  be a bijective continuous function. If  $X$  is compact and  $Y$  is Hausdorff, then  $f$  is a homeomorphism.*

*Proof.* We shall prove that images of closed sets of  $X$  under  $f$  are closed in  $Y$ ; this will prove continuity of the map  $f^{-1}$ . If  $A$  is closed in  $X$ , then  $A$  is compact, by Theorem 26.2. Therefore, by the theorem just proved,  $f(A)$  is compact. Since  $Y$  is Hausdorff,  $f(A)$  is closed in  $Y$ , by Theorem 26.3. ■

**Theorem 26.7.** *The product of finitely many compact spaces is compact.*

*Proof.* We shall prove that the product of two compact spaces is compact; the theorem follows by induction for any finite product.

*Step 1.* Suppose that we are given spaces  $X$  and  $Y$ , with  $Y$  compact. Suppose that  $x_0$  is a point of  $X$ , and  $N$  is an open set of  $X \times Y$  containing the “slice”  $x_0 \times Y$  of  $X \times Y$ . We prove the following:

*There is a neighborhood  $W$  of  $x_0$  in  $X$  such that  $N$  contains the entire set  $W \times Y$ .*

The set  $W \times Y$  is often called a **tube** about  $x_0 \times Y$ .

First let us cover  $x_0 \times Y$  by basis elements  $U \times V$  (for the topology of  $X \times Y$ ) lying in  $N$ . The space  $x_0 \times Y$  is compact, being homeomorphic to  $Y$ . Therefore, we can cover  $x_0 \times Y$  by finitely many such basis elements

$$U_1 \times V_1, \dots, U_n \times V_n.$$

(We assume that each of the basis elements  $U_i \times V_i$  actually intersects  $x_0 \times Y$ , since otherwise that basis element would be superfluous; we could discard it from the finite collection and still have a covering of  $x_0 \times Y$ .) Define

$$W = U_1 \cap \dots \cap U_n.$$

The set  $W$  is open, and it contains  $x_0$  because each set  $U_i \times V_i$  intersects  $x_0 \times Y$ .

We assert that the sets  $U_i \times V_i$ , which were chosen to cover the slice  $x_0 \times Y$ , actually cover the tube  $W \times Y$ . Let  $x \times y$  be a point of  $W \times Y$ . Consider the point  $x_0 \times y$  of the slice  $x_0 \times Y$  having the same  $y$ -coordinate as this point. Now  $x_0 \times y$  belongs to  $U_i \times V_i$  for some  $i$ , so that  $y \in V_i$ . But  $x \in U_j$  for every  $j$  (because  $x \in W$ ). Therefore, we have  $x \times y \in U_i \times V_i$ , as desired.

Since all the sets  $U_i \times V_i$  lie in  $N$ , and since they cover  $W \times Y$ , the tube  $W \times Y$  lies in  $N$  also. See Figure 26.2.

*Step 2.* Now we prove the theorem. Let  $X$  and  $Y$  be compact spaces. Let  $\mathcal{A}$  be an open covering of  $X \times Y$ . Given  $x_0 \in X$ , the slice  $x_0 \times Y$  is compact and may therefore be covered by finitely many elements  $A_1, \dots, A_m$  of  $\mathcal{A}$ . Their union  $N = A_1 \cup \dots \cup A_m$  is an open set containing  $x_0 \times Y$ ; by Step 1, the open set  $N$  contains




Figure 26.2

a tube  $W \times Y$  about  $x_0 \times Y$ , where  $W$  is open in  $X$ . Then  $W \times Y$  is covered by finitely many elements  $A_1, \dots, A_m$  of  $\mathcal{A}$ .

Thus, for each  $x$  in  $X$ , we can choose a neighborhood  $W_x$  of  $x$  such that the tube  $W_x \times Y$  can be covered by finitely many elements of  $\mathcal{A}$ . The collection of all the neighborhoods  $W_x$  is an open covering of  $X$ ; therefore by compactness of  $X$ , there exists a finite subcollection

$$\{W_1, \dots, W_k\}$$

covering  $X$ . The union of the tubes

$$W_1 \times Y, \dots, W_k \times Y$$

is all of  $X \times Y$ ; since each may be covered by finitely many elements of  $\mathcal{A}$ , so may  $X \times Y$  be covered. ■

The statement proved in Step 1 of the preceding proof will be useful to us later, so we repeat it here as a lemma, for reference purposes:

**Lemma 26.8 (The tube lemma).** *Consider the product space  $X \times Y$ , where  $Y$  is compact. If  $N$  is an open set of  $X \times Y$  containing the slice  $x_0 \times Y$  of  $X \times Y$ , then  $N$  contains some tube  $W \times Y$  about  $x_0 \times Y$ , where  $W$  is a neighborhood of  $x_0$  in  $X$ .*

EXAMPLE 7. The tube lemma is certainly not true if  $Y$  is not compact. For example, let  $Y$  be the  $y$ -axis in  $\mathbb{R}^2$ , and let

$$N = \{x \times y; |x| < 1/(y^2 + 1)\}.$$

Then  $N$  is an open set containing the set  $0 \times \mathbb{R}$ , but it contains no tube about  $0 \times \mathbb{R}$ . It is illustrated in Figure 26.3.